

AECV90-R4

Growth, Feeding and Fecundity of Fish in Gleniffer Lake (Dickson Dam Reservoir)

CANADIANA

C 2

MAR 26 1991



Alberta
ENVIRONMENTAL CENTRE



Digitized by the Internet Archive
in 2016

GROWTH, FEEDING AND FECUNDITY OF FISH IN
GLENIFFER LAKE (DICKSON DAM RESERVOIR)

by

James W. Moore

Aquatic Biology Branch

Alberta Environmental Centre

Vegreville, Alberta

August 29, 1990

GROWTH, FEEDING AND FECUNDITY OF FISH IN
GLENIFFER LAKE (DICKSON DAM RESERVOIR)

JAMES W. MOORE
FISHERY BIOLOGY BRANCH
ALBERTA ENVIRONMENTAL CENTRE
VEGREVILLE, ALBERTA

MOORE, J.W. 1990

This report may be cited as:

Moore, J.W. 1990. Growth, Feeding and Fecundity of Fish in Gleniffer Lake (Dickson Dam Reservoir). Alberta Environmental Centre, Vegreville, AB. AECV90-R4. 40 pp.

TABLE OF CONTENTS

| | Page |
|---|------|
| LIST OF TABLES | iii |
| LIST OF FIGURES | v |
| SUMMARY | vii |
| PROJECT TEAM | ix |
| ACKNOWLEDGMENTS | ix |
| 1. INTRODUCTION | 1 |
| 1.1 Background | 1 |
| 1.2 Purpose of Study | 1 |
| 2. STUDY AREA | 4 |
| 2.1 Red Deer River | 4 |
| 2.2 History of Dam and Reservoir Facility | 4 |
| 2.3 History of Fish Plantings | 5 |
| 3. MATERIALS AND METHODS | 5 |
| 3.1 Fish Collections | 5 |
| 3.1.1 Beach Seine | 5 |
| 3.1.2 Gill Net | 9 |
| 3.2 Aging Methods | 10 |
| 3.3 Stomach Content Analysis | 11 |
| 3.4 Fecundity Analysis | 11 |
| 3.5 Statistical Methods | 12 |
| 4. RESULTS | 12 |
| 4.1 Length/Weight Relation | 12 |
| 4.2 Length/Age Relation | 19 |

TABLE OF CONTENTS

| | Page |
|--------------------------------|------|
| 4.3 Stomach Contents | 27 |
| 4.3.1 Mountain Whitefish | 27 |
| 4.3.2 Rainbow Trout | 27 |
| 4.3.3 Northern Pike | 28 |
| 4.3.4 Burbot | 28 |
| 4.4 Fecundity | 28 |
| 5. DISCUSSION | 32 |
| REFERENCES | 39 |

LIST OF TABLES

| | |
|--|----|
| Table 1. Summary of information on Dickson Dam, Gleniffer Lake and surrounding area | 6 |
| Table 2. Summary of fish plantings carried out at Gleniffer Lake .. | 7 |
| Table 3. Description of beach seine sites | 7 |
| Table 4. Aging structures used in this study | 9 |
| Table 5. Regression parameters relating wet weight (g) (dependent variable) to fork length (mm) (independent variable) in fish caught by gill net from Gleniffer Lake during 1984-1987 | 13 |
| Table 6. t-test of regression slopes relating age to weight for fish caught during 1984, 1985, 1986 and 1987 | 19 |

LIST OF TABLES (continued)

Page

| | |
|---|----|
| Table 7. Regression parameters relating fork length (mm) (independent variable) to age (years) (dependent variable) in fish caught by gill net from Gleniffer Lake during 1984-1987 | 20 |
| Table 8. t-test of regression slopes relating age to fork length for fish caught during 1984, 1985, 1986 and 1987 | 21 |
| Table 9. Fork length (mm) of fish of standard age (years) collected by gill net from Gleniffer Lake during 1984-1987 | 27 |
| Table 10. Summary of stomach content analyses | 29 |
| Table 11. Consumption of fish by northern pike and burbot in Gleniffer Lake 1984-87 | 30 |
| Table 12. Regression parameters relating length (mm) to calculated total number of eggs in fish caught by gill net from Gleniffer Lake during 1984-1987 | 32 |
| Table 13. Mean fork length (mm) of mountain whitefish at age 3+ and 6+ years in various lakes and rivers | 33 |
| Table 14. Mean fork length (mm) of rainbow trout at age 3+ and 6+ years in various lakes | 34 |
| Table 15. Mean fork length (mm) of northern pike at age 3+ and 6+ years in various Alberta lakes | 36 |
| Table 16. Mean fork length (mm) of burbot at age 3+ and 6+ years in various lakes | 37 |
| Table 17. Mean fork length (mm) of white sucker at age 3+ and 6+ years in various lakes | 38 |

LIST OF TABLES (continued)

Page

| | |
|---|----|
| Table 18. Mean fork length (mm) of longnose sucker at age 3+ and 6+ years in various lakes and rivers | 38 |
|---|----|

LIST OF FIGURES

| | |
|---|----|
| Figure 1. Location of Gleniffer Lake | 2 |
| Figure 2. Beach seine sampling sites | 8 |
| Figure 3. Gill net sampling sites | 8 |
| Figure 4. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of mountain whitefish and rainbow trout collected by gill net from Gleniffer Lake during 1984-87 | 14 |
| Figure 5. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of northern pike collected by gill net from Gleniffer Lake during 1984-87 | 15 |
| Figure 6. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of burbot collected by gill net from Gleniffer Lake during 1984-87 | 16 |
| Figure 7. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of longnose sucker collected by gill net from Gleniffer Lake during 1984-87 | 17 |

LIST OF FIGURES

Page

| | |
|---|----|
| Figure 8. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of white sucker collected by gill net from Gleniffer Lake during 1984-87 | 18 |
| Figure 9. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of mountain whitefish and rainbow trout collected by gill net from Gleniffer Lake during 1984-87 | 22 |
| Figure 10. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of northern pike collected by gill net from Gleniffer Lake during 1984-87 | 23 |
| Figure 11. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of burbot collected by gill net from Gleniffer Lake during 1984-87 | 24 |
| Figure 12. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of white sucker collected by gill net from Gleniffer Lake during 1984-87 | 25 |
| Figure 13. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of longnose sucker collected by gill net from Gleniffer Lake during 1984-87 | 26 |
| Figure 14. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to calculated total number of eggs of burbot, northern pike, rainbow trout, longnose sucker and white sucker collected by gill net from Gleniffer Lake during 1984-87 | 31 |

SUMMARY

Fish were collected by gill net and beach seine from a newly formed reservoir (Gleniffer Lake, Alberta) during a four-year period (1984-1987). The reservoir is located west of the City of Red Deer on the Red Deer River. Collections were made in the reservoir at standard sites and times throughout the study. Length, weight and age were determined for mountain whitefish (Prosopium williamsoni), northern pike (Esox lucius), burbot (Lota lota), white sucker (Catostomus commersoni), and longnose sucker (Catostomus catostomus). In addition, stomach content analysis was completed on mountain whitefish, rainbow trout, northern pike and burbot; fecundity was determined for rainbow trout (Oncorhynchus mykiss), northern pike, burbot, white sucker and longnose sucker. From these collections it was observed that:

- i) mountain whitefish were initially trapped by the impoundment of the Red Deer River, but either died out or emigrated from the reservoir by 1985;
- ii) growth of mountain whitefish and rainbow trout from the reservoir was greater than that observed for populations in many other Alberta lakes;
- iii) based on back-calculated growth rates, young (<3 years old) northern pike apparently grew better in the Red Deer River than Gleniffer Lake whereas older pike (6+ years old) grew better in the reservoir than the river; this may have

(vii)

reflected a temporary reduction in food for young pike in the first years after impoundment;

- iv) although growth of burbot was largely unaffected by impoundment, reduced growth of young burbot was noted during 1985;
- v) white sucker and longnose sucker grew significantly faster in the reservoir than the Red Deer River;
- vi) although rainbow trout were planted in large numbers in the reservoir, they did not form a significant part of the diet of predacious species such as northern pike and burbot;
- vii) the diet of mountain whitefish and rainbow trout was typical of that reported elsewhere in Canada and the USA;
- viii) fecundity of rainbow trout, northern pike, burbot, white sucker, and longnose sucker was typical of that reported for other northern temperate zone lakes.

PROJECT TEAM

| | |
|-----------------|----------------|
| Brenda Dew | James Moore |
| Karen Blumhagen | Ed Paleczny |
| Bernie Goski | David Park |
| John Craig | Dan Patton |
| Sandra Gellings | Jane Schneider |
| Lee George | James Somers |
| Darrel Katan | Kevin Smiley |

ACKNOWLEDGMENTS

We are indebted to staff from Fish and Wildlife Division (particularly D. Lowe, Head, Fisheries Management, Red Deer Region, and T. Mill, Director, Fisheries Management) for their assistance during this study. In addition, staff at the dam facility (T.J. Maduke, Reservoir Manager and his staff) provided essential logistic support for this study.

We are grateful to M. Barrett and Z. Florence for their assistance in reviewing and calculating much of the statistical data in this report.

We also wish to recognize M. Herbut for his assistance with the graphic presentation of data required for this report.

1. INTRODUCTION

1.1 Background

Gleniffer Lake, located on the Red Deer River west of the City of Red Deer, was formed with the completion of the Dickson Dam in 1983 (Figure 1). The reservoir has an area of 1734 ha at full storage and measures approximately 11 km in length and 2 km in width. The facility provides two primary benefits: i) an assured water supply for downstream communities, and ii) improved water quality downstream.

During the spring and summer, runoff water is captured in the reservoir. Enough water is released during the winter to ensure a minimum flow of at least $16 \text{ m}^3 \text{ s}^{-1}$. This rate of flow meets the present needs of downstream industries and communities including the Cities of Red Deer and Drumheller, and allows for future growth.

1.2 Purpose of Study

Because of its size and accessibility, the reservoir has the potential to become a significant sport fishing facility. The purpose of this report is to describe the growth, feeding and fecundity of major fish species in the reservoir. Such information is useful in managing the fisheries and, in particular, determining the nature and extent of fish plantings, and the relative success of different fish species.

A previous report (Alberta Environmental Centre, 1989a) concerning fish in the reservoir indicated that:




Figure 1. Location of Gleniffer Lake.

- i) the most frequently captured species in the reservoir were white sucker (Catostomus commersoni), longnose sucker (Catostomus catostomus), northern pike (Esox lucius), and burbot (Lota lota). Although rainbow trout (Oncorhynchus mykiss) had been planted in large numbers in the reservoir, they made up only a small part of the catch;
- ii) the catch per unit effort (CUE) using gill nets was low, ranging from 1 - 4.7 fish (all species) per 1000 m net per h;
- iii) planted rainbow trout and brown trout (Salmo trutta) apparently failed to spawn in the reservoir;
- iv) the populations of northern pike, white sucker, longnose sucker and burbot apparently expanded immediately after impoundment, and later decreased in numbers;
- v) the most frequently captured fish by anglers were rainbow trout, brown trout and northern pike;
- vi) the CUE for angling was extremely low; depending on year, 11-45 h were required to angle one fish.

Another report (Alberta Environmental Centre, 1989b) concerning mercury in fish from the reservoir indicated that:

- i) residues were low, generally $<500 \mu\text{g kg}^{-1}$ (0.5 ppm) in all species;
- ii) there was no increase in tissue residues as the reservoir aged;
- iii) no limits needed to be placed on consumption of fish from Gleniffer Lake.

2. STUDY AREA

2.1 Red Deer River

The Red Deer River originates in Banff National Park at an elevation of approximately 1830 m above sea level. The river flows 185 km through the Rocky Mountains to Gleniffer Lake which is at an elevation of 948 m above sea level at Full Storage Level (FSL). The drainage area to this point is 5520 km². The river continues a further 571 km eastward across parkland and prairies where it joins the South Saskatchewan River near Alberta's border with Saskatchewan.

2.2 History of Dam and Reservoir Facility

Technical studies began in 1971 and, after five years of investigations and a series of public hearings, a decision was made to build the dam. Clearing of the site and construction of access roads began in February 1980. By the spring of 1983, the dam was completed and the reservoir began filling.

The service spillway can handle any major flood of a magnitude occurring once in 10,000 years. In the unlikely event of a larger flood, an emergency spillway could be used to divert excess flow. A flood forecasting system using remote rainfall gauges and satellite communications gives early warning to operators about oncoming floods. The diversion tunnels are equipped for installation of hydroelectric power facilities, if needed. A summary of information on the dam and reservoir is listed in Table 1.

2.3 History of Fish Plantings

Rainbow trout and cutthroat trout (Salmo clarki) were first planted in Gleniffer Lake in 1983 (Table 2). The introductions of rainbow trout continued through 1987, but no further cutthroat trout were planted beyond 1983. Brown trout were introduced to the reservoir only during 1984. Total plantings for all species exceeded one million fish.

3. MATERIALS AND METHODS

3.1 Fish Collections

Fish used in this study were collected with gill nets and beach seine during 1984, 1985, 1986 and 1987.

3.1.1 Beach Seine

The seine net was 18 m in length, 1.8 m in depth with a center bag of 1.8 m x 1.8 m. The mesh was 6.4 mm delta white nylon except for the bag which had a mesh of 3.2 mm.

Four sites in each of three basins (west, central and east) were selected for a total of 12 sites (Figure 2, Table 3). These sites were sampled once a month from May through September.

After capture, fish were given a lethal dose of the anaesthetic MS222, weighed and measured, and then preserved in 10% formalin for later identification and enumeration. Stomach contents and gonads were preserved in a 10% formalin solution.

Table 1. Summary of information on Dickson Dam, Gleniffer Lake and surrounding area.

| | | |
|--|--------------------------------------|--|
| | DAM | |
| Crest Length | 650 m | |
| Height | 40 m | |
| Volume of Fill | 3,600,000 m ³ | |
| Crest Elevation | 952 m | |
| | TUNNELS (2) | |
| Length | 525 m | |
| Diameter | 5.5 m | |
| Design Discharge | 38 m ³ s ⁻¹ | |
| | SPILLWAY | |
| Weir Elevation | 940.5 m | |
| Weir Length | 60 m | |
| Structure Length | 235 m | |
| Vertical Gates (5) | 8.6 x 9.6 m | |
| Design Discharge | 2,600 m ³ s ⁻¹ | |
| | EMERGENCY SPILLWAY | |
| Channel Width | 130 m | |
| Design Discharge | 2,800 m ³ s ⁻¹ | |
| | RESERVOIR | |
| Flooded Area (Full Storage Level) | 1,734 ha | |
| Usable Storage | 203 10 ⁶ m ³ | |
| Reservoir Length | 11 km | |
| Reservoir Width | 2 km | |
| Full Storage Level | 948 m elevation | |
| Annual Flush Rate | 5.5 x | |
| Proportion of Incoming Sediment Retained | 88 % | |
| Annual Amount of Sediment Retained | 290,000 m ³ | |
| Life Expectancy | 500 yrs | |
| | DYKES | |
| Length (North Side) | 3.0 km | |
| Length (South Side) | 3.7 km | |
| | DRAINAGE BASIN | |
| Area | 5,520 km ² | |
| Mean Run-off | 32 m ³ yr ⁻¹ | |

Table 2. Summary of fish plantings carried out at Gleniffer Lake.

| Year | Species | Month(s) Planted | Number Planted | Length (cm) |
|--------|---------------------|-------------------|----------------|-------------|
| 1983 | Rainbow Trout | Sept. | 61,557 | 4.8- 20 |
| 1983 | Cutthroat Trout | Sept. | 135,635 | ~ 5.5 |
| 1984 | Rainbow Trout | June, Aug., Sept. | 152,600 | 6.4 - >20 |
| 1984 | Brown Trout | Aug., Sept. | 67,100 | 5.2 - >20 |
| 1985 | Rainbow Trout | May, June, Aug. | 498,500 | 5.6 - 12 |
| 1986 | Rainbow Trout | Aug. | 180,00 | >15 |
| 1986 | Rainbow Trout | Sept., Oct. | 375 | >20 |
| 1987 | Rainbow Trout | Aug. | 242 | >20 |
| TOTALS | Rainbow Trout | | 893,274 | 4.8- >20 |
| 83-87 | Cutthroat Trout | | 135,635 | 5.5 |
| | Brown Trout | | 67,100 | 5.2- >20 |
| GRAND | | | | |
| TOTAL | Trout (all species) | | 1,096,009 | 4.8 - >20 |

SOURCE: Fish Planting Lists (1983-1987) issued by Alberta Forestry, Lands and Wildlife.

Table 3. Description of beach seine sites.

| Location (see Fig. 2) | Maximum Depth (m) | Description of Bottom Texture |
|--------------------------|----------------------|----------------------------------|
| East A | 2 | Cobble, sand |
| East B | 2 | Soft sand/clay |
| East C | 2 | Gravel, soft sand/clay |
| East D | 7.5 | Cobble |
| Central A | 2.5 | Sand/gravel |
| Central B | 3 | Sand |
| Central C | 2 | Soft sand/clay |
| Central D | 4.5 | Cobble, some sand |
| West A | 4 | Sand |
| West B | 2.5 | Sand |
| West C | 3 | Cobble, sand |
| West D | 4 | Sand |




Figure 3. Gill net sampling sites.




Figure 4. Seine sampling sites. Deep set (>10 m); shallow set (<10 m).

3.1.2 Gill Net

Multi-panel gill nets, measuring 50 m in length, were used in all collections. The nets, 1.8 m deep, were made of green or colourless monofilament nylon. Two nets were usually tied together, yielding a total length of 100 m. Mesh sizes were 1.9, 2.5, 3.8, 5.1, 5.4, 7.6, 8.9, 10.2, 11.4 and 12.7 cm.

There were two sampling sites in each of the three basins (Figure 3). One site was situated in shallow (<10 m) water and the other in deep (>10 m) water. Selection of an actual site was based on its suitability for gill netting, specifically minimum slope and minimum debris content.

Each net was set in the morning and left for 24 h. The fish were taken to a field laboratory where the species, fork length, and wet weight of each fish were determined. Condition factor ($\text{Weight}^3/\text{Fork Length}$) was calculated. Stomach contents and ovaries were preserved in a 10% formalin solution. Aging structures were also taken (Table 4).

Table 4. Aging structures used in this study.

| Species | Structure | Year |
|--------------------|------------------------|--------------------------|
| Mountain Whitefish | Fin ray | 1984 |
| Rainbow Trout | Fin ray | 1986 |
| Burbot | Otolith | 1984, 1985, 1986 |
| Northern Pike | Cleithrum Operculum | 1985, 1986, 1987 1984 |
| White Sucker | Fin ray Scales | 1984, 1986, 1987 1985 |
| Longnose Sucker | Fin ray | All years |

3.2 Aging Methods

All aging structures were placed in labelled envelopes. In the laboratory, the envelopes and structures were frozen at -20°C until preparation. The preparation procedures used prior to aging were:

Scales

- Five scales were selected at random.
- The scales were cleaned with water and/or mildly basic solution and placed between two glass slides.
- The sides were taped, labelled and stored in a slide tray.

Fin Rays

- Two fin rays were air dried for 2–5 days.
- The proximal end of each ray was coated with epoxy and allowed to set for 24 h.
- The ray was cut into 1-mm sections using a jeweller's saw.
- The sections were mounted with Diatex mounting fluid on a glass slide.
- The slides were labelled and stored in a slide tray.

Cleithra and Opercula

- One of each structure was first cleaned with warm water and soft brushes, then air dried and allowed to clear.
- Both structures were returned to the labelled scale envelope before evaluation.

Otoliths

- Two otoliths were removed from each fish.
- Larger otoliths were sectioned and allowed to air dry. Small otoliths were air dried without sectioning.
- All structures were returned to the labelled envelope before evaluation.

Ages were determined for each specimen through the enumeration of annuli.

3.3 Stomach Content Analysis

The field-preserved stomach samples of rainbow trout, northern pike, mountain whitefish and burbot were sorted in the laboratory and identified according to five taxonomic classifications. Ingested fish were identified to species whenever possible. After identification, the samples were dried at 55°C to constant weight. Stomachs of other species were not examined due to time and financial restraints.

3.4 Fecundity Analysis

The ovary was sampled in six areas (2 x posterior, 2 x middle, 2 x anterior). A total of 1000 eggs (250 for rainbow trout) was counted from each of the six samples. The enumerated eggs were then dried to constant weight. The remainder of the ovary was also dried, thereby permitting an estimate to be made of total egg numbers.

Because mountain whitefish were caught only in 1984, at a time prior to sexual maturation, no determination of fecundity could be made of that species.

3.5 Statistical Analysis

The following relationships were expressed as linear regressions: fork length-weight, fork length-age, fork length-number of eggs. Linear regression was used because it provided the best fit for the majority of data. The regression slopes relating fork length to age and fork length to weight were analyzed using the t-test.

4. RESULTS

4.1 Length/Weight Relation

The species collected in sufficient numbers to warrant analysis were northern pike, white sucker, longnose sucker, mountain whitefish, burbot, and rainbow trout. Mountain whitefish were collected only during 1984 whereas rainbow trout were collected in sufficient numbers to warrant analysis during 1986 and 1987. Only 3 brown trout, 1 brook trout (Salvelinus fontinalis) and 1 dolly varden (Salvelinus malma) and no cutthroat trout were caught during the entire study.

The linear regressions relating wet weight to fork length of the above-noted species are listed in Table 5 and graphed in Figures 4-8. The slopes of these regressions were analyzed using t-test (Table 6). In the case of rainbow trout and northern pike, there were no significant differences among slopes during the study. This means that the length-weight relationship remained constant for both species in all years. Burbot, on the other hand, carried more weight in fish of standard length caught in 1986 and 1987 compared to 1985. This can possibly be related to improved feeding conditions as the reservoir


Table 5. Regression parameters relating wet weight (g) (dependent variable) to fork length (mm) (independent variable) in fish caught by gill net from Gleniffer Lake during 1984-1987.

| Year | Slope | Y-intercept | r ² | Mean Condition Factor* | N |
|--------------------|-------|-------------|----------------|------------------------|-----|
| MOUNTAIN WHITEFISH | | | | | |
| 1984 | 1.991 | -318 | 0.89 | 1.06 | 24 |
| RAINBOW TROUT | | | | | |
| 1986 | 13.38 | -4893 | 0.96 | 1.33 | 11 |
| 1987 | 8.15 | -2236 | 0.60 | 1.43 | 6 |
| NORTHERN PIKE | | | | | |
| 1984 | 6.063 | -1905 | 0.59 | 0.89 | 23 |
| 1985 | 8.201 | -3018 | 0.90 | 0.82 | 35 |
| 1986 | 7.685 | -2740 | 0.86 | 0.77 | 20 |
| 1987 | 6.946 | -2148 | 0.85 | 0.76 | 25 |
| BURBOT | | | | | |
| 1984 | 3.366 | -861 | 0.89 | 0.61 | 11 |
| 1985 | 4.595 | -1441 | 0.91 | 0.57 | 10 |
| 1986 | 5.472 | -1796 | 0.90 | 0.59 | 13 |
| 1987 | 6.968 | -2870 | 0.97 | 0.50 | 6 |
| WHITE SUCKER | | | | | |
| 1984 | 5.069 | -1110 | 0.91 | 1.47 | 31 |
| 1985 | 3.831 | -694 | 0.94 | 1.37 | 54 |
| 1986 | 4.939 | -1087 | 0.93 | 1.39 | 50 |
| 1987 | 4.959 | -1078 | 0.94 | 1.44 | 67 |
| LONGNOSE SUCKER | | | | | |
| 1984 | 3.876 | -762 | 0.91 | 1.35 | 56 |
| 1985 | 5.376 | -1304 | 0.88 | 1.32 | 117 |
| 1986 | 6.161 | -1621 | 0.84 | 1.30 | 116 |
| 1987 | 5.048 | -1168 | 0.93 | 1.27 | 80 |


* $\text{Weight}^3/\text{Length}$

MOUNTAIN WHITEFISH

1984

**RAINBOW TROUT**

1986



1987




Figure 4. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of mountain whitefish and rainbow trout collected by gill net from Gleniffer Lake during 1984-87.




Figure 5. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of northern pike collected by gill net from Gleniffer Lake during 1984-87.




Figure 6. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of burbot collected by gill net from Gleniffer Lake during 1984-87.




Figure 7. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of longnose sucker collected by gill net from Gleniffer Lake during 1984-87.




Figure 8. Linear regression analysis ($\pm 95\%$ confidence limits) relating weight to length of white sucker collected by gill net from Gleniffer Lake during 1984-87.

Table 6. t-test of regression slopes relating age to weight for fish caught during 1984, 1985, 1986, and 1987.

| Species/Year | 1985 | 1986 | 1987 |
|-------------------------|------|------|------|
| Rainbow Trout 1986 | | | NS |
| Northern Pike 1984 | NS | NS | NS |
| 1985 | | NS | NS |
| 1986 | | | NS |
| Burbot 1984 | NS | S | S |
| 1985 | | NS | S |
| 1986 | | | S |
| White Sucker 1984 | S | NS | NS |
| 1985 | | S | S |
| 1986 | | | NS |
| Longnose Sucker 1984 | S | S | S |
| 1985 | | S | NS |
| 1985 | | | S |

S = significant at $P=0.05$ NS= not significant at $P=0.05$

aged (see Discussion). Although the same situation applied to longnose sucker, the relationship between weight and length in white sucker did not change in 1984, 1986 and 1987.

4.2 Length/Age Relation

The regression slopes relating age and fork length for fish collected in different years were analyzed using t-test (Tables 7 and 8; Figures 9-13). In the case of longnose sucker, white sucker and northern pike, fish at standard age had a greater fork length in

Table 7. Regression parameters relating fork length (mm) (independent variable) to age (years) (dependent variable) in fish caught by gill net from Gleniffer Lake during 1984-1987.

| Year | Slope | Y-intercept | r^2 | N |
|--------------------|-------|-------------|-------|-----|
| MOUNTAIN WHITEFISH | | | | |
| 1984 | 22.4 | 153 | 0.45 | 24 |
| RAINBOW TROUT | | | | |
| 1986 | 73.2 | 134 | 0.42 | 11 |
| 1987 | -6.6 | 553 | 0.12 | 6 |
| NORTHERN PIKE | | | | |
| 1984 | 13.4 | 446 | 0.05 | 23 |
| 1985 | 37.7 | 372 | 0.67 | 35 |
| 1986 | 61.7 | 237 | 0.70 | 20 |
| 1987 | 68.4 | 170 | 0.94 | 25 |
| BURBOT | | | | |
| 1984 | 31.1 | 274 | 0.65 | 11 |
| 1985 | 70.5 | 107 | 0.88 | 10 |
| 1986 | 45.5 | 208 | 0.31 | 13 |
| 1987 | 68.8 | 106 | 0.79 | 6 |
| WHITE SUCKER | | | | |
| 1984 | 31.8 | 217 | 0.61 | 31 |
| 1985 | 82.9 | 89 | 0.72 | 54 |
| 1986 | 71.9 | 89 | 0.54 | 50 |
| 1987 | 53.2 | 135 | 0.74 | 67 |
| LONGNOSE SUCKER | | | | |
| 1984 | 26.1 | 228 | 0.37 | 56 |
| 1985 | 31.0 | 247 | 0.44 | 117 |
| 1986 | 10.5 | 353 | 0.08 | 116 |
| 1987 | 50.1 | 115 | 0.59 | 80 |

Table 8. t-test of regression slopes relating age to fork length for fish caught during 1984, 1985, 1986 and 1987.

| Species/Year | 1985 | 1986 | 1987 |
|-------------------------|------|------|------|
| Rainbow Trout 1986 | | | S |
| Northern Pike 1984 | NS | S | S |
| 1985 | | NS | S |
| 1986 | | | NS |
| Burbot 1984 | S | NS | NS |
| 1985 | | NS | NS |
| 1986 | | | NS |
| White Sucker 1984 | S | S | S |
| 1985 | | NS | NS |
| 1986 | | | NS |
| Longnose Sucker 1984 | NS | S | S |
| 1985 | | NS | S |
| 1986 | | | S |

S = significant at $P=0.05$.


NS= not significant at $P=0.05$.

1987 than in 1984. Burbot, on the other hand, showed no significant change in the length/age relationship between 1984 and 1987.

The length/age relationship for rainbow trout was significantly different between 1986 and 1987, presumably reflecting the introduction of hatchery-reared fish of different length. For example, trout aged 3+ years averaged 354 mm in length in 1986, increasing to 523 mm in 1987 (Table 9). However, trout decreased in length (from 523 to 513 mm) between ages 3+ and 6+.

MOUNTAIN WHITEFISH

1984



1986

RAINBOW TROUT

1987





Figure 9. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of mountain whitefish and rainbow trout collected by gill net from Gleniffer Lake during 1984-87.




Figure 10. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of northern pike collected by gill net from Gleniffer Lake during 1984-87.




Figure 11. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of burbot collected by gill net from Gleniffer Lake during 1984-87.




Figure 12. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of white sucker collected by gill net from Gleniffer Lake during 1984-87.




Figure 13. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to age of longnose sucker collected by gill net from Gleniffer Lake during 1984-87.

Table 9. Fork length (mm) of fish of standard age (years) collected by gill net from Gleniffer Lake during 1984-1987.

| Year | Standard Age | | | | | | | | | | | |
|------|--------------------|-----|---------------|-----|---------------|-----|--------|-----|--------------|-----|-----------------|-----|
| | 3+ | 6+ | 3+ | 6+ | 3+ | 6+ | 3+ | 6+ | 3+ | 6+ | 3+ | 6+ |
| | Mountain Whitefish | | Rainbow Trout | | Northern Pike | | Burbot | | White Sucker | | Longnose Sucker | |
| 1984 | 220 | 287 | ND | ND | 486 | 526 | 367 | 461 | 312 | 408 | 306 | 384 |
| 1985 | ND | ND | ND | ND | 485 | 598 | 318 | 530 | 338 | 586 | 340 | 433 |
| 1986 | ND | ND | 354 | 573 | 422 | 607 | 344 | 481 | 305 | 520 | 385 | 417 |
| 1987 | ND | ND | 523 | 513 | 375 | 580 | 312 | 520 | 295 | 454 | 265 | 415 |

ND = no data

4.3 Stomach Contents

4.3.1 Mountain Whitefish

This species, collected only in 1984, fed mainly on cladocerans (*Daphnia* sp.) and insects (Table 10). The cladoceran component accounted for 95-99% of the dry weight of stomach contents, which ranged from 0.03 to 0.31 g dry weight per fish.

4.3.2 Rainbow Trout

Although rainbow trout were captured in all years, detailed stomach analysis was limited to the 1986 and 1987 collections, which included a relatively large number of specimens. In both years, insects were present in the stomachs of most fish (Table 10), whereas

plant material and unidentified remains were also found in many stomachs. The total weight of stomach contents ranged from 0.01 g to 5.5 g dry weight per fish.

4.3.3 Northern Pike

The majority of stomachs of northern pike collected in 1985 and 1986 were empty (Tables 10 and 11). In 1984 and 1987, however, fish and unidentified remains were found in the majority of stomachs. The most frequently consumed fish were stickleback (Gasterosteus sp.), sucker (unidentified) and mountain whitefish (Prosopium williamsoni) (only in 1984) (Tables 10 and 11). Only one rainbow trout was found in the pike stomachs. Weight of stomach contents ranged from 0.0 g to 20.4 g dry weight per fish.

4.3.4 Burbot

Burbot were also piscivorous - approximately 45% of all stomach examined contained fish (Tables 10 and 11). The most frequently ingested species was stickleback (unidentified). Insects were also numerically important (Table 10) but, in terms of weight, formed <1% of ingested material (Table 10). Weight of stomach contents ranged from 0.0 g to 5.4 g dry weight per fish.

4.4 Fecundity

Based on the regression parameters in Table 12, burbot of standard fork length (600 mm) contained approximately 774×10^3 eggs/fish.

Table 10. Summary of stomach content analyses.

| Year | Number of Fish | Number of Stomachs Containing Item | | | | |
|--------------------|----------------|------------------------------------|---------|-------------|----------|----------------------|
| | | Fish | Insects | Cladocerans | Molluscs | Unidentified Remains |
| MOUNTAIN WHITEFISH | | | | | | |
| 1984 | 45 | 0 | 25 | 31 | 0 | 12 |
| RAINBOW TROUT | | | | | | |
| 1986 | 17 | 0 | 12 | 1 | 1 | 8 |
| 1987 | 8 | 0 | 8 | 1 | 4 | 7 |
| NORTHERN PIKE | | | | | | |
| 1984 | 25 | 13 | 4 | 1 | 0 | 6 |
| 1985 | 35 | 8 | 0 | 0 | 0 | 0 |
| 1986 | 20 | 2 | 0 | 0 | 0 | 0 |
| 1987 | 25 | 11 | 0 | 0 | 0 | 10 |
| BURBOT | | | | | | |
| 1984 | 12 | 5 | 5 | 0 | 1 | 2 |
| 1985 | 11 | 5 | 4 | 1 | 0 | 3 |
| 1986 | 13 | 6 | 5 | 0 | 0 | 5 |
| 1987 | 6 | 3 | 2 | 0 | 1 | 1 |

Northern pike of the same length contained 48×10^3 eggs/fish. The corresponding number of eggs/fish for rainbow trout (500 mm), longnose sucker (400 mm) and white sucker (400 mm) is 4300, 32,200 and 43,700 respectively. In all species, the number of eggs increased with the length of the fish (Figure 14).

Table 11. Consumption of fish by northern pike and burbot in Gleniffer Lake 1984-87.

| Year | Number of Fish | Number of Stomachs Containing Fish | Number and Species of Ingested Fish | | | | | |
|---------------|----------------|------------------------------------|-------------------------------------|---------------|--------------------------|-------------------------------|--------|--------------|
| | | | Mountain Whitefish | Rainbow Trout | Sucker (species unknown) | Stickleback (species unknown) | Burbot | Unidentified |
| | | | | | | | | |
| NORTHERN PIKE | | | | | | | | |
| 1984 | 25 | 13 | 11 | 1 | 1 | 34 | 0 | 12 |
| 1985 | 35 | 8 | 0 | 0 | 5 | 0 | 1 | 5 |
| 1986 | 20 | 2 | 0 | 0 | 2 | 0 | 0 | 0 |
| 1987 | 25 | 11 | 1 | 0 | 12 | 0 | 0 | 9 |
| BURBOT | | | | | | | | |
| 1984 | 12 | 5 | 0 | 0 | 0 | 13 | 0 | 14 |
| 1985 | 11 | 5 | 0 | 0 | 0 | 29 | 1 | 14 |
| 1986 | 13 | 6 | 0 | 0 | 0 | 29 | 1 | 14 |
| 1987 | 6 | 3 | 0 | 0 | 0 | 0 | 0 | 3 |




Figure 14. Linear regression analysis ($\pm 95\%$ confidence limits) relating length to calculated total number of eggs of burbot, northern pike, rainbow trout, longnose sucker and white sucker collected by gill net from Gleniffer Lake during 1984-87.

Table 12. Regression parameters relating length (mm) to calculated total number of eggs in fish caught by gill net from Gleniffer Lake during 1984-1987.

| Species | Slope | Y-intercept | r^2 | N |
|-----------------|-------|-------------|-------|----|
| Rainbow Trout | 24.9 | -8,154 | 0.46 | 13 |
| Northern Pike | 248.9 | -101,500 | 0.76 | 41 |
| Burbot | 3250 | -1,176,100 | 0.84 | 10 |
| White Sucker | 145.9 | -14,649 | 0.29 | 51 |
| Longnose Sucker | 259.0 | -71,449 | 0.42 | 42 |

5. DISCUSSION

Mountain whitefish apparently were trapped by impoundment of the Red Deer River in 1983. Although inhabiting the reservoir for another year, the population either died out or emigrated from the reservoir by 1985. This is typical of whitefish populations in other parts of Canada. For example, Bodaly et al. (1984) attributed the post-impoundment collapse of the lake whitefish (Coregonus clupeaformis) to emigration from the reservoir. Similarly, increased sedimentation in reservoirs reduces the rate of survival of lake whitefish eggs (Fudge and Bodaly, 1984).

Because mountain whitefish could have spent no more than 1 year in the reservoir prior to capture in 1984, most of their growth was due to life in the river rather than the reservoir. These whitefish were

relatively large compared to lake-dwelling fish, but were slightly smaller than those from the North Saskatchewan River (Table 13). No comparative records appear to be available regarding growth of mountain whitefish in the Red Deer River upstream of the reservoir.

Because rainbow trout of different age and size were planted in Gleniffer Lake, reliable determinations of growth after release cannot be made. In 1986, for example, the average fork length of 3+ years trout was 343 mm whereas, in 1987, the corresponding length was 523 mm. This reflects the planting of relatively large hatchery reared fish in the 1984-year class. In comparison to other lakes, the

Table 13. Mean fork length (mm) of mountain whitefish at age 3+ and 6+ years in various lakes and rivers.

| Lake/River | Age (Years) | | Reference |
|------------------------------|-------------|-----|------------|
| | 3+ | 6+ | |
| Gleniffer Lake, AB | 220 | 287 | This study |
| Bow Lake, AB | 104 | 142 | 1 |
| Pyramid Lake, AB | 150 | 268 | 2 |
| Kananaskis Lake, AB | 171 | 267 | 1 |
| Waterton Lake, AB | 171 | 260 | 1 |
| Lake Minnewanka, AB | 171 | 273 | 1 |
| Ghost River Reservoir, AB | 199 | NR | 1 |
| Cultus Lake, BC | 218 | 305 | 1 |
| North Saskatchewan River, AB | 247 | 334 | 3 |

SOURCES: 1. McHugh (1941)
 2. Rawson and Elsey (1948)
 3. Alberta Forestry, Lands and Wildlife (1989)
 NR = not reported

trout planted in Gleniffer Lake were relatively large at age 3+ years (Table 14), but this reflects a combination of growth in the hatchery and reservoir.

Northern pike which were 3+ years old and caught in 1984 must have been spawned in 1981 and presumably spent the first 2-3 years of their life in the Red Deer River. On the other hand, pike of the same age

Table 14. Mean fork length (mm) of rainbow trout at age 3+ and 6+ years in various lakes.

| Lake | Age (Years) | | Reference |
|----------------------|-------------|-----|------------|
| | 3+ | 6+ | |
| Gleniffer, AB (1986) | 354 | 573 | This study |
| Gleniffer, AB (1987) | 523 | 513 | This study |
| Pyramid, AB | 190 | 376 | 1 |
| Okanagan, BC | 290 | NR | 1 |
| Kootenay, BC | 303 | 590 | 1 |
| Watch, BC | 291 | NR | 1 |
| Cluculz, BC | 275 | NR | 1 |
| Lightning, BC | 183 | NR | 1 |
| Pennask, BC | 203 | NR | 1 |
| Loon, BC | 213 | NR | 1 |
| Bouche, BC | 222 | NR | 1 |
| Big Bar, BC | 247 | NR | 1 |
| Kelly, BC | 252 | NR | 1 |
| Beaver, BC | 257 | NR | 1 |
| Dairy, BC | 354 | NR | 1 |
| Knouff, BC | 356 | NR | 1 |
| Pavilion, BC | 380 | NR | 1 |
| Glimpse, BC | 387 | NR | 1 |
| Peterhope, BC | 468 | NR | 1 |

SOURCE: 1. Larkin et al. (1956)

NR = not reported

captured in 1987 were spawned in 1984 and had presumably spent all of their life in the reservoir. From Table 15, it is apparent that the 1987-caught fish were much smaller than the 1984-caught fish, implying that the river provided better habitat for the growth of young pike than the reservoir. The specific factors influencing the difference in growth are not known at this time. From age 3+ to 6+ years, pike grew much faster in the reservoir than the river (Table 15). It is possible that impoundment of the river significantly curtailed the production of fry and other potential food fish for young pike. As the reservoir matured, however, reproduction of potential food species increased. Rainbow trout, introduced in large numbers to the reservoir, was not a significant food of northern pike.

Fork length of northern pike at age 6+ years was comparable to fish from many other Alberta lakes (Table 15). For comparison the age 3+ pike caught in 1987 from the reservoir were shorter than fish from 10 other lakes listed in Table 15. Once again, the factors influencing the differences in growth are not known.

The growth of burbot was largely unaffected by impoundment, with the apparent exception of 1984-1985. During that period, the regression slopes relating length to age changed significantly: 3+ year old fish caught in 1985 were smaller than those caught in 1984 while 6+ year old burbot were larger in 1985 than 1984. As with northern pike, it appears that the river provided better habitat for the growth of young burbot whereas the reservoir was better for the growth of older burbot. This trend was not apparent in subsequent

Table 15. Mean fork length (mm) of northern pike at age 3+ and 6+ years in various Alberta lakes.

| Lake | Age (Years) | | Reference |
|------------------|-------------|-----|------------|
| | 3+ | 6+ | |
| Gleniffer (1984) | 486 | 526 | This study |
| Gleniffer (1985) | 485 | 598 | This study |
| Gleniffer (1986) | 422 | 607 | This study |
| Gleniffer (1987) | 375 | 580 | This study |
| Amisk | 431 | NR | 1 |
| Beaver | 488 | 561 | 1 |
| Cold | 463 | 635 | 1 |
| Elinor | 411 | 485 | 1 |
| Garner | 508 | 571 | 1 |
| Lac La Biche | 476 | 666 | 1 |
| Lac Ste. Anne | 385 | NR | 1 |
| Muriel | 490 | 595 | 1 |
| Pinehurst | 464 | 601 | 1 |
| Touchwood | 431 | 591 | 1 |

SOURCE: 1. Alberta Forestry, Lands and Wildlife (1983-1986)
NR = not reported

years (Table 7). The overall growth of burbot in the reservoir was intermediate between populations in northern and more temperate lakes (Table 16).

White sucker and longnose sucker grew significantly faster in the reservoir than in the river (Tables 7, 8). The older fish were most affected whereas young suckers showed no increase in growth. As with northern pike, it appears that the absence of a good food base during the first years of impoundment limited growth. Suckers feed primarily on benthic invertebrates, a group of organisms that would be virtually

Table 16. Mean fork length (mm) of burbot at age 3+ and 6+ years in various lakes.

| Lake | Age (Years) | | Reference |
|----------------------|-------------|-----|------------|
| | 3+ | 6+ | |
| Gleniffer, AB (1984) | 367 | 461 | This study |
| Gleniffer, AB (1985) | 318 | 530 | This study |
| Gleniffer, AB (1986) | 344 | 481 | This study |
| Gleniffer, AB (1987) | 312 | 520 | This study |
| Heming, MN | 279 | 399 | 1 |
| Superior, WI | 300 | 439 | 2 |
| Erie, ON | 376 | 540 | 1 |
| Simcoe, ON | 432 | 572 | 1 |

SOURCES: 1. Scott and Crossman (1973)
 2. Becker (1983)

eliminated during reservoir construction but which could rapidly recolonize the substrate. This reflects the ability of many invertebrate species such as insects to complete their life cycle in <1 year and to move long distances. Sucker growth in the reservoir was relatively rapid, even when compared to populations in more temperate climates (Tables 17, 18).

Since only one rainbow trout was found in the stomachs of 34 northern pike from the reservoir, it appears that the rate of predation is low. Similarly, no trout were found in the stomachs of burbot (42 specimens examined). This means that the success of the planting program for the reservoir was probably not diminished by the presence of predacious species.

Table 17. Mean fork length (mm) of white sucker at age 3+ and 6+ years in various lakes.

| Lake | Age (Years) | | Reference |
|-----------------------|-------------|------------------|------------|
| | 3+ | 6+ | |
| Gleniffer, AB (1984) | 312 | 408 | This study |
| Gleniffer, AB (1985) | 338 | 586 | This study |
| Gleniffer, AB (1986) | 305 | 520 | This study |
| Gleniffer, AB (1987) | 295 | 454 | This study |
| Waskiv, SK | 120 | 300 | 1 |
| Muskellunge, WI | 163 | 262 | 2 |
| Lake of the Woods, ON | 284 | 396 | 1 |
| George, ON | 336 | 452 | 1 |
| Michigan, WI | 362 | 430 | 2 |
| Winnebago, WI | 406 | 498 ^a | 2 |

SOURCES: 1. Scott and Crossman (1973)
2. Becker (1983)

^a females only

Table 18. Mean fork length (mm) of longnose sucker at age 3+ and 6+ years in various lakes and rivers.

| Lake/River | Age (Years) | | Reference |
|------------------------------------|-------------|-----|------------|
| | 3+ | 6+ | |
| Gleniffer, AB (1984) | 306 | 384 | This study |
| Gleniffer, AB (1985) | 340 | 433 | This study |
| Gleniffer, AB (1986) | 385 | 417 | This study |
| Gleniffer, AB (1987) | 265 | 415 | This study |
| Pyramid, AB | 107 | 178 | 1 |
| Great Slave Lake (northern), NWT | 170 | 314 | 1 |
| Great Slave Lake (whole lake), NWT | 177 | 288 | 2 |
| Great Slave Lake (southern), NWT | 203 | 308 | 1 |
| North Saskatchewan River, SK | 323 | 439 | 1 |

SOURCES: 1. Scott and Crossman (1973)
2. Harris (1962)

The diet of the other species examined in this study is typical of that reported elsewhere in Canada and the USA (Scott and Crossman, 1973; Becker, 1983).

REFERENCES

- Alberta Environmental Centre. 1989a. Species composition and angler use of fish in Gleniffer Lake (Dickson Dam reservoir). Alberta Environmental Centre, Vegreville, AB. AECV89-R5. 38 pp.
- Alberta Environmental Centre. 1989b. A five-year study of mercury in fish from a newly formed reservoir (Gleniffer Lake, Alberta). Alberta Environmental Centre, Vegreville, AB. AECV89-R4. 24 pp.
- Alberta Forestry, Lands and Wildlife. 1983-1986. Unpublished data. Fish and Wildlife Division, St. Paul, AB.
- Alberta Forestry, Lands and Wildlife. 1989. Unpublished data. Fish and Wildlife Division, Edmonton, AB.
- Becker, G. 1983. Fishes of Wisconsin. University of Wisconsin Press. Madison, WI. 1052 pp.
- Bodaly, R.A., T.W.D. Johnson, R.J.P. Fudge, and J.W. Clayton. 1984. Collapse of the lake whitefish (Coregonus clupeaformis) fishery in Southern Indian Lake, Manitoba, following lake impoundment and river diversion. Can. J. Fish. Aquat. Sci. 41:692-700.
- Fish Planting Lists. 1983-1987. Fish and Wildlife Division, Alberta Forestry, Lands and Wildlife, Edmonton, AB.

- Fudge, R.J.P., and R.A. Bodaly. 1984. Postimpoundment winter sedimentation and survival of lake whitefish (Coregonus clupeaformis) eggs in Southern Indian Lake, Manitoba. Can. J. Fish. Aquat. Sci. 41:701-705.
- Harris, Roy H.D. 1962. Growth and reproduction of the longnose sucker, Catostomus catostomus, in Great Slave Lake. J. Fish. Res. Bd. Can. 19:113-126.
- Larkin, P.A., J.G. Terpenning, and R.R. Parker. 1956. Size as a determinant of growth rate in rainbow trout. Trans. Am. Fish. Soc. 86:84-96.
- McHugh, J.L. 1941. Growth of the Rocky Mountain whitefish. J. Fish. Res. Bd. Can. 5:337-343.
- Rawson, D.S., and C.A. Elsey. 1948. Reduction in the longnose sucker population of Pyramid Lake, Alberta, in an attempt to improve angling. Trans. Amer. Fish. Soc. 78:13-31.
- Scott, W.B., and E.J. Crossman. 1973. Freshwater fishes of Canada. Fish. Res. Bd. Can. Bull. 184:641-645.

N.L.C. - B.N.C.



3 3286 10458965 6